

Acta Crystallographica Section C

**Crystal Structure  
Communications**

ISSN 0108-2701

Editor: **Anthony Linden**


## The crucial role of C—H···O and C=O··· $\pi$ interactions in the building of three-dimensional structures of dicarboxylic acid–biimidazole compounds

**Xiao-Li Gao, Li-Ping Lu and Miao-Li Zhu***Acta Cryst.* (2009). **C65**, o123–o127

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site or institutional repository provided that this cover page is retained. Republication of this article or its storage in electronic databases other than as specified above is not permitted without prior permission in writing from the IUCr.

For further information see <http://journals.iucr.org/services/authorrights.html>



*Acta Crystallographica Section C: Crystal Structure Communications* specializes in the rapid dissemination of high-quality studies of crystal and molecular structures of interest in fields such as chemistry, biochemistry, mineralogy, pharmacology, physics and materials science. The numerical and text descriptions of each structure are submitted to the journal electronically as a Crystallographic Information File (CIF) and are checked and typeset automatically prior to peer review. The journal is well known for its high standards of structural reliability and presentation. *Section C* publishes approximately 1000 structures per year; readers have access to an archive that includes high-quality structural data for over 10000 compounds.

Crystallography Journals **Online** is available from [journals.iucr.org](http://journals.iucr.org)

# The crucial role of C—H···O and C=O··· $\pi$ interactions in the building of three-dimensional structures of dicarboxylic acid–biimidazole compounds

Xiao-Li Gao, Li-Ping Lu\* and Miao-Li Zhu\*

Institute of Molecular Science, Key Laboratory of Chemical Biology and Molecular Engineering of the Education Ministry, Shanxi University, Taiyuan, Shanxi 030006, People's Republic of China

Correspondence e-mail: luliping@sxu.edu.cn, miaoli@sxu.edu.cn

Received 15 January 2009

Accepted 18 February 2009


Online 7 March 2009

The supramolecular architectures of three dicarboxylic acid–biimidazole compounds, namely, 2,2'-biimidazolium malonate,  $C_6H_8N_4^{2+} \cdot C_3H_2O_4^{2-}$ , (I), 2,2'-bi(1*H*-imidazole) succinic acid,  $C_6H_6N_4 \cdot C_4H_6O_4$ , (II), and 2,2'-biimidazolium 2,2'-iminodiacetate chloride,  $C_6H_8N_4^{2+} \cdot C_4H_6NO_4^- \cdot Cl^-$ , (III), are reported. The crystal structures are assembled by the same process, namely double conventional N—H···O or O—H···N hydrogen bonds link the dicarboxylates and biimidazoles to form tapes, which are stacked in parallel through lone-pair–aromatic interactions between carbonyl O atoms and biimidazole groups and are further linked *via* weak C—H···O interactions. The C=O··· $\pi$  interactions involved in stacking the tapes in (II) and the C—H···O interactions involved in linking the tapes in (II) and (III) demonstrate the crucial role of these interactions in the crystal packing. There is crystallographically imposed symmetry in all three structures. In (I), two independent malonate anions have their central C atoms on twofold axes and two biimidazolium dication each lie about independent inversion centres; in (II), the components lie about inversion centres, while in (III), the unique cation lies about an inversion centre and the iminodiacetate and chloride anions lie across and on a mirror plane, respectively.


## Comment

Noncovalent weak interactions, such as C—H···O and C=O··· $\pi$ , have attracted much interest. These weak interactions have been widely discussed in relation to the crystal packing of organic molecules and the determination of the folded structures of biological molecules (Derewenda *et al.*, 1995; Desiraju, 1996, 2005; Khurram *et al.*, 2006; Jain *et al.*, 2007; Lu *et al.*, 2007; Wan *et al.*, 2008). However, self-assembled supramolecular architectures are often stabilized as a

result of the synergy of a variety of weak interactions (Khurram *et al.*, 2006; Shukla *et al.*, 2007; Wan *et al.*, 2008). It is difficult to distinguish the effect of an individual weak interaction. Desiraju (2005) suggested that weak interactions can be classified as innocuous, supportive or intrusive. We describe here the decisive role of carbonyl– $\pi$  and C—H···O interactions in the assembly of the supramolecular architectures of three dicarboxylic acid–biimidazole compounds, (I)–(III).




The asymmetric unit of (I) contains two independent malonate anions, with their central C atoms on twofold axes, and two biimidazolium cations, each lying about independent inversion centres (Fig. 1). Strong N—H···O hydrogen bonds




**Figure 1**

A view of the structure of (I), with displacement ellipsoids drawn at the 30% probability level. Dashed lines represent hydrogen bonds. [Symmetry codes: (i)  $-x + 1, -y, -z + 1$ ; (ii)  $-x, -y, -z + 1$ ; (iv)  $-x, y, -z + \frac{1}{2}$ ; (ix)  $-x + 1, y, -z + \frac{1}{2}$ .]


**Figure 2**


The structure built from N—H...O hydrogen bonds and C—H...O interactions in (I). Dashed lines represent hydrogen bonds. [Symmetry codes: (ii)  $-x, -y, -z + 1$ ; (v)  $x, -y + 1, z + \frac{1}{2}$ ; (vi)  $-x + 1, -y + 1, -z + 1$ ; (viii)  $x, y + 1, z$ ; (ix)  $-x + 1, y + 1, -z + \frac{3}{2}$ .]


**Figure 3**

The C=O... $\pi$  interaction (double dashed lines) between neighbouring A tapes and B tapes in (I). Dotted lines represent hydrogen bonds. [Symmetry codes: (v)  $x, -y + 1, z + \frac{1}{2}$ ; (vi)  $-x + 1, -y + 1, -z + 1$ ; (viii)  $x, y + 1, z$ ; (x)  $-x, y + 1, -z + \frac{3}{2}$ ; (xi)  $-x, y, -z + \frac{3}{2}$ .]

(Fig. 1 and Table 1) link the malonate and biimidazolium moieties to form two distinct units, which further assemble into two different zigzag tapes designated as A and B, as shown as Fig. 2. Four C—H...O interactions (Fig. 2 and Table 1) link the two tapes into a three-dimensional network, and concomitantly there are three possible C=O... $\pi$ (imidazole ring) interactions, one of which takes part in the stacking of tapes A while the other two link tapes B, as described in Table 1 and shown in Fig. 3.


In (II), there is crystallographically imposed inversion symmetry with the succinic acid and neutral biimidazole molecules lying about inversion centres. The succinic acid and


**Figure 4**


A view of the structure of (II), with displacement ellipsoids drawn at the 30% probability level. Dashed lines represent hydrogen bonds. [Symmetry codes: (i)  $-x + 2, -y + 1, -z + 1$ ; (iv)  $-x, -y + 1, -z$ .]

biimidazole molecules are linked by pairs of O—H...N and N—H...O hydrogen bonds (Fig. 4 and Table 2), leading to linear tapes which are further linked to form sheets in the (10 $\bar{2}$ ) plane by the C3—H3...O2( $x + 1, -y + \frac{1}{2}, z + \frac{1}{2}$ ) interaction (Fig. 5 and Table 2). The sheets assemble into a three-dimensional structure by the C4=O1... $\pi$ (imidazole ring) ( $-x + 1, -y + 1, -z + 1$ ) interaction (Fig. 6 and Table 2).

In (III) (Fig. 7), the biimidazolium dication lies about an inversion centre, the iminodiacetate ion lies across a mirror



**Figure 5**

Part of the two-dimensional sheet assembled from O—H...N, N—H...O and C—H...O interactions in (II). Dashed lines represent hydrogen bonds. [Symmetry codes: (i)  $-x + 2, -y + 1, -z + 1$ ; (ii)  $x + 1, -y + \frac{1}{2}, z + \frac{1}{2}$ ; (v)  $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$ .]


**Figure 6**

The  $C=O \cdots \pi$  interaction (double dashed lines) between neighbouring sheets in (II). Dotted lines represent hydrogen bonds. [Symmetry codes: (iii)  $-x + 1, -y + 1, -z + 1$ ; (vi)  $x + 2, y, z + 1$ ; (vii)  $-x + 3, -y + 1, -z + 2$ .]


plane and the chloride ion lies on a mirror plane. Iminodiacetate anions and biimidazolium dication are linked to form wave-like tapes by pairs of  $N-H \cdots O$  hydrogen bonds (Fig. 8 and Table 3). These tapes are linked to form sheets by weak  $C3-H3 \cdots O1(-x + 1, -y + 1, -z)$  interactions (Fig. 8 and Table 3). The sheets are packed to form a three-dimensional network *via* two  $N-H \cdots Cl$  hydrogen bonds and two  $C5=O1 \cdots \pi$  interactions to imidazole rings at  $(-x, -y + 1, -z)$  and  $(-x + 1, -y + 1, -z + 1)$  (Fig. 9 and Table 3).

**Figure 7**

A view of the structure of (III), with displacement ellipsoids drawn at the 30% probability level. Dashed lines represent hydrogen bonds. [Symmetry codes: (i)  $-x, -y + 1, -z + 1$ ; (vii)  $x, -y + \frac{3}{2}, z$ .]

**Figure 8**

The two-dimensional sheet assembled from  $N-H \cdots O$  and  $C-H \cdots O$  interactions in (III). Dashed lines represent hydrogen bonds. [Symmetry codes: (i)  $-x, -y + 1, -z + 1$ ; (iii)  $-x + 1, y - \frac{1}{2}, -z + 1$ ; (iv)  $-x + 1, -y + 1, -z$ ; (viii)  $x, y, z - 1$ .]

**Figure 9**

The  $C=O \cdots \pi$  interactions (double dashed lines) between neighbouring sheets in (III). Dotted lines represent hydrogen bonds. [Symmetry codes: (vii)  $x, -y + \frac{3}{2}, z$ ; (ix)  $-x, y + \frac{3}{2}, -z + 1$ ; (x)  $x + 1, y, z + 1$ ; (xi)  $x + 1, -y + \frac{3}{2}, z + 1$ ; (xii)  $-x + 1, -y, -z + 1$ .]

The supramolecular structures of (I), (II) and (III) reveal that they are assembled by the same process, namely that pairs of  $N-H \cdots O$  or  $O-H \cdots N$  hydrogen bonds link the dicarboxylate and biimidazole molecules to form tapes, which are stacked in parallel through lone-pair-aromatic interactions between carbonyl O atoms and biimidazole groups and are further linked *via* weak  $C-H \cdots O$  interactions.

## Experimental

Diimidazole (1 mmol) and malonic, succinic or iminodiacetic acid (1 mmol) were dissolved in water (10 ml) by adding 0.7–0.9 ml of 2 M HCl while stirring. The solutions were left to stand at room temperature and colourless crystals of (I) and (II) and yellow crystals of (III) were obtained after several days.

## Compound (I)

### Crystal data

$C_6H_8N_4^{2+} \cdot C_3H_2O_4^{2-}$   
 $M_r = 238.21$   
 Monoclinic,  $P2_1/c$   
 $a = 15.663(5) \text{ \AA}$   
 $b = 4.4319(14) \text{ \AA}$   
 $c = 18.221(4) \text{ \AA}$   
 $\beta = 124.517(18)^\circ$

$V = 1042.2(5) \text{ \AA}^3$   
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.12 \text{ mm}^{-1}$   
 $T = 298 \text{ K}$   
 $0.40 \times 0.40 \times 0.40 \text{ mm}$

Data collection

Bruker SMART 1K CCD area-detector diffractometer  
Absorption correction: multi-scan (SADABS; Sheldrick, 2000)  
 $T_{\min} = 0.845$ ,  $T_{\max} = 0.953$

4027 measured reflections  
1841 independent reflections  
1519 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.014$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$   
 $wR(F^2) = 0.122$   
 $S = 1.07$   
1841 reflections

155 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.28 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.23 \text{ e } \text{\AA}^{-3}$

Compound (II)

Crystal data

$\text{C}_6\text{H}_6\text{N}_4 \cdot \text{C}_4\text{H}_6\text{O}_4$   
 $M_r = 252.24$   
Monoclinic,  $P2_1/c$   
 $a = 4.906 (3) \text{ \AA}$   
 $b = 13.887 (8) \text{ \AA}$   
 $c = 8.468 (5) \text{ \AA}$   
 $\beta = 94.839 (8)^\circ$

$V = 574.9 (6) \text{ \AA}^3$   
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.12 \text{ mm}^{-1}$   
 $T = 298 \text{ K}$   
 $0.30 \times 0.30 \times 0.27 \text{ mm}$

Data collection

Bruker SMART 1K CCD area-detector diffractometer  
Absorption correction: multi-scan (SADABS; Sheldrick, 2000)  
 $T_{\min} = 0.845$ ,  $T_{\max} = 0.970$

2390 measured reflections  
1055 independent reflections  
868 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$   
 $wR(F^2) = 0.117$   
 $S = 1.05$   
1055 reflections

82 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.19 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.15 \text{ e } \text{\AA}^{-3}$

Compound (III)

Crystal data

$\text{C}_6\text{H}_8\text{N}_4^{2+} \cdot \text{C}_4\text{H}_6\text{NO}_4^- \cdot \text{Cl}^-$   
 $M_r = 303.71$   
Monoclinic,  $P2_1/m$   
 $a = 5.3095 (13) \text{ \AA}$   
 $b = 22.941 (6) \text{ \AA}$   
 $c = 5.7023 (14) \text{ \AA}$   
 $\beta = 107.930 (3)^\circ$

$V = 660.8 (3) \text{ \AA}^3$   
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.31 \text{ mm}^{-1}$   
 $T = 298 \text{ K}$   
 $0.40 \times 0.40 \times 0.40 \text{ mm}$

Data collection

Bruker SMART 1K CCD area-detector diffractometer  
Absorption correction: multi-scan (SADABS; Sheldrick, 2000)  
 $T_{\min} = 0.747$ ,  $T_{\max} = 0.886$

3079 measured reflections  
1158 independent reflections  
1027 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.017$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.064$   
 $wR(F^2) = 0.180$   
 $S = 1.05$   
1158 reflections

94 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.69 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.32 \text{ e } \text{\AA}^{-3}$

For compound (III), the systematic absences permitted  $P2_1$  or  $P2_1/m$  as possible space groups;  $P2_1/m$  was selected and confirmed by the structure analysis. H atoms attached to C atoms were placed in geometrically idealized positions and refined with  $U_{\text{iso}}(\text{H})$  values of  $1.2U_{\text{eq}}(\text{C})$ . H atoms attached to N and O atoms were located from

Table 1

Hydrogen-bond and contact geometry ( $\text{\AA}$ ,  $^\circ$ ) for (I).

$\text{Cg1}$  and  $\text{Cg2}$  are the centroids of the  $\text{N1/C1/N2/C3/C2}$  and  $\text{N3/C4/N4/C6/C5}$  rings, respectively.

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N1}-\text{H1} \cdots \text{O3}^{\text{i}}$	0.86	1.78	2.612 (2)	163
$\text{N2}-\text{H2A} \cdots \text{O4}$	0.86	1.80	2.646 (2)	166
$\text{N3}-\text{H3A} \cdots \text{O1}^{\text{ii}}$	0.86	1.77	2.609 (2)	166
$\text{N4}-\text{H4} \cdots \text{O2}$	0.86	1.79	2.626 (2)	164
$\text{C2}-\text{H2} \cdots \text{O2}^{\text{iii}}$	0.93	2.58	3.401 (2)	148
$\text{C3}-\text{H3} \cdots \text{O1}^{\text{iv}}$	0.93	2.41	3.292 (2)	158
$\text{C5}-\text{H5} \cdots \text{O4}^{\text{v}}$	0.93	2.70	3.317 (2)	125
$\text{C6}-\text{H6} \cdots \text{O3}^{\text{vi}}$	0.93	2.45	3.345 (2)	162
$\text{C7}-\text{O1} \cdots \text{Cg2}^{\text{vii}}$	1.25 (1)	3.49 (1)	3.714 (3)	90
$\text{C9}-\text{O3} \cdots \text{Cg1}^{\text{viii}}$	1.26 (1)	3.54 (1)	3.796 (2)	92
$\text{C9}-\text{O4} \cdots \text{Cg1}^{\text{viii}}$	1.24 (1)	3.68 (1)	4.538 (3)	127

Symmetry codes: (i)  $-x+1, -y, -z+1$ ; (ii)  $-x, -y, -z+1$ ; (iii)  $x, y-1, z$ ; (iv)  $-x, y, -z+\frac{1}{2}$ ; (v)  $x, -y+1, z+\frac{1}{2}$ ; (vi)  $-x+1, -y+1, -z+1$ ; (vii)  $-x, -y+1, -z+1$ ; (viii)  $x, y+1, z$ .

Table 2

Hydrogen-bond and contact geometry ( $\text{\AA}$ ,  $^\circ$ ) for (II).

$\text{Cg1}$  is the centroid of the  $\text{N1/C1/N2/C3/C2}$  ring.

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O2}-\text{H2} \cdots \text{N1}$	0.85	1.74	2.575 (2)	169
$\text{N2}-\text{H2B} \cdots \text{O1}^{\text{i}}$	0.86	1.93	2.779 (2)	170
$\text{C3}-\text{H3} \cdots \text{O2}^{\text{ii}}$	0.93	2.41	3.309 (3)	162
$\text{C4}-\text{O1} \cdots \text{Cg1}^{\text{iii}}$	1.22 (1)	3.37 (1)	3.720 (3)	97

Symmetry codes: (i)  $-x+2, -y+1, -z+1$ ; (ii)  $x+1, -y+\frac{1}{2}, z+\frac{1}{2}$ ; (iii)  $-x+1, -y+1, -z+1$ .

Table 3

Hydrogen-bond and contact geometry ( $\text{\AA}$ ,  $^\circ$ ) for (III).

$\text{Cg1}$  is the centroid of the  $\text{N1/C1/N2/C2/C3}$  ring.

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N1}-\text{H1} \cdots \text{O1}$	0.82	1.79	2.592 (3)	167
$\text{N2}-\text{H2A} \cdots \text{O2}^{\text{i}}$	0.82	1.83	2.643 (3)	171
$\text{N3}-\text{H3A} \cdots \text{Cl1}$	0.90	2.56	3.271 (7)	137
$\text{N3}-\text{H3B} \cdots \text{Cl1}^{\text{ii}}$	0.90	2.50	3.244 (7)	141
$\text{C2}-\text{H2} \cdots \text{Cl1}^{\text{iii}}$	0.93	2.63	3.520 (3)	161
$\text{C3}-\text{H3} \cdots \text{O1}^{\text{iv}}$	0.93	2.38	3.253 (4)	156
$\text{C5}-\text{O1} \cdots \text{Cg1}^{\text{v}}$	1.26 (1)	3.33 (1)	3.816 (4)	103
$\text{C5}-\text{O1} \cdots \text{Cg1}^{\text{vi}}$	1.26 (1)	3.50 (1)	4.029 (4)	106

Symmetry codes: (i)  $-x, -y+1, -z+1$ ; (ii)  $x-1, y, z-1$ ; (iii)  $-x+1, y-\frac{1}{2}, -z+1$ ; (iv)  $-x+1, -y+1, -z$ ; (v)  $-x, -y+1, -z$ ; (vi)  $-x+1, -y+1, -z+1$ .

difference Fourier maps and refined using a riding model, with  $U_{\text{iso}}(\text{H})$  values of  $1.2U_{\text{eq}}(\text{N})$  or  $1.5U_{\text{eq}}(\text{O})$  of their parent atoms.

For all compounds, data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL/PC* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL/PC* and *ORTEP-3* (Farrugia, 1997).

The authors acknowledge the financial support of the National Natural Science Foundation of China (grant No.

20471033) and the Natural Science Foundation of Shanxi Province of China (grant No. 20051013), as well as the Overseas Returned Scholar Foundation of Shanxi Province of China in 2006 and 2008.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: FG3076). Services for accessing these data are described at the back of the journal.

## References

- Bruker (2000). *SMART* (Version 5.0) and *SAINT* (Version 6.02). Bruker AXS Inc., Madison, Wisconsin, USA.
- Derewenda, Z. S., Lee, L. & Derewenda, U. (1995). *J. Mol. Biol.* **252**, 248–262.
- Desiraju, G. R. (1996). *Acc. Chem. Res.* **29**, 441–449.
- Desiraju, G. R. (2005). *Chem. Commun.* pp. 2995–3001.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Jain, A., Purohit, C. S., Verma, S. & Sankararamakrishnan, R. (2007). *J. Phys. Chem. B*, **111**, 8680–8683.
- Khurram, M., Qureshi, N. & Smith, M. D. (2006). *Chem. Commun.* pp. 5006–5008.
- Lu, Z., Gamez, P., Mutikainen, L., Turpeinen, U. & Reedijk, J. (2007). *Cryst. Growth Des.* **7**, 1669–1671.
- Sheldrick, G. M. (2000). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shukla, R., Lindeman, S. V. & Rathore, R. (2007). *Chem. Commun.* pp. 3717–3719.
- Wan, C.-Q., Chen, X.-D. & Mak, T. C. W. (2008). *CrystEngComm*, **10**, 475–478.